

A Data Quality Screening Service for Remote Sensing Data



Christopher Lynnes, NASA/GSFC

Edward Olsen, NASA/JPL

Peter Fox, RPI

Bruce Vollmer, NASA/GSFC

Robert Wolfe, NASA/GSFC

Contributions from: N. Most, Y.-I. Won, T. Hearty, R. Strub, S. Ahmad, S. Zednik, M. Hegde and A. Rezaiyan-Nojani

Funded by: NASA's Advancing Collaborative Connections for Earth System Science (ACCESS) Program



Outline



- Why a Data Quality Screening Service?
- Making Quality Information Easier to Use via the Data Quality Screening Service (DQSS)
- Ruture Directions for DQSS



Why a Data Quality Screening Service?

The quality of AIRS data varies considerably




			Do Not Use (%)	
Total Precipitable Water	38	38	24	
Carbon Monoxide	64	7	29	
Surface Temperature	5	44	51	


Version 5 Level 2 Standard Retrieval Statistics



Quality schemes can be relatively simple...











...or they can be more complicated



Hurricane Ike, viewed by the Atmospheric Infrared Sounder (AIRS)



Air Temperature at 300 mbar




PBest: Maximum pressure for which quality value is "Best" in temperature profiles



Quality flags are also sometimes packed together into bytes





Bitfield arrangement for the Cloud_Mask_SDS variable in atmospheric products from Moderate Resolution Imaging Spectroradiometer (MODIS)



Current user scenarios...



- - Search for and download data
 - Compare the compared of the compared
 - Read & understand documentation on quality
 - Write custom routine to filter out bad pixels

Repeat for each user


- Equally likely scenario (especially in user communities not familiar with satellite data)
 - Search for and download data
 - Assume that quality has a negligible effect





The effect of bad quality data is often not negligible



Hurricane Ike, 9/10/2008










Neglecting quality may introduce bias (a more subtle effect)



AIRS Relative Humidity Comparison against Dropsonde with and without Applying PBest Quality Flag Filtering Boxed data points indicate AIRS RH data with dry bias > 20%



From a study by Sun Wong (JPL) on specific humidity in the Atlantic Main Development Region for Tropical Storms



Making Quality Information Easier to Use via the Data Quality Screening Service (DQSS)

•



The DQSS filters out bad pixels for the user




- © Default user scenario
 - Search for data
 - Select science team recommendation for quality screening (filtering)
 - □ Download screened data
- More advanced scenario
 - Search for data
 - Select custom quality screening parameters
 - □ Download screened data




DQSS replaces bad-quality pixels with fill values




Original data array (Total column precipitable water)



Mask based on user criteria (Quality level < 2)



Good quality data pixels retained




Output file has the same format and structure as the input file (except for extra mask and original_data fields)




Visualizations help users see the effect of different quality filters

Data within product file



Best quality only

Best + Good quality





DQSS can encode the science team recommendations on quality screening



- AIRS Level 2 Standard Product
 - Use Best-only for data assimilation uses
 - Use Best+Good for climatic studies
- R MODIS Aerosols

 - Use Marginal+Good+VeryGood over ocean



Or, users can select their own criteria...

0	Goddard Earth Sciences (GES) Data and Information Center (DISC)							
▶ 🙆 A A 🥞 -	► 🛣 A A 👣 + 🚭 http://mirador.gsfc.nasa.gov/cgi-bin/mirador/serviceSelection.pl?CGISESSID=9989335638b24c60469170d6fc250273&Selecter & Q Google							
AIRX2F	I settings are based on Science Tea (Note: "Good" retains retrievals that ou can choose settings for all para	t Good or better).						
(Sele	Apply To All Parameters: ct Good to screen all parameters following Science Team criteria)	☐ Best * Good **						
* For data assimilation, use of Best quality data only is recommended. ** Using only Good quality (or better) data is recommended for most uses.								
+ News + Restricted Data	or parameter by	y parameter.						
	Temperature_Parameters							
	TSurfStd Visualize	☐ Best ⓒ Good ☐ NoScreening						
	TAirStd Visualize	○ Best Good NoScreening						
	TAirMWOnlyStd Visualize	Best Good NoScreening						
	TSurfAir Visualize	Best Good NoScreening						
Definition Demonstration								



Status and Future





DQSS Status



- DQSS is operational for AIRS L2 Standard Products
 - DQSS is offered through the Mirador data search interface at the GES DISC
- Usage Metrics will be collected:
 - Real Basic usage
 - What criteria are used
 - Screening invocation is a simple URL GET
 - What the "yield" was for the screening (future)



Future: Extend DQSS to Other Datasets



- Moderate Resolution Imaging Spectroradiometer (MODIS) L2 Atmospheres
- Microwave Limb Sounder (MLS) L2
- © Ozone Monitoring Instrument (OMI) L2
- Resolution Dynamics Limb Sounder (HIRDLS)



Combine DQSS with Other Services



™ Subsetting

Reformatting to NetCDF

○ OPeNDAP

Suggestions?



DQSS Target Uses and Users



	Routine	Visual- ization	Quick Recon.	Machine- level	Metrics
Interdisciplinary	X	X			
Educational	X	X			
Expert/Power	?		X	X	
Applications				X	
Algorithm Developers					X



DQSS Recap



- Screening satellite data can be difficult and time consuming for users
- The Data Quality Screening System will provide an easy-to-use service
- The result should be:
 - More attention to quality on users' part
 - More accurate handling of quality information...